

Were these symptoms present on most days for at least one month?*
Kentucky BRFSS 2000

Geographic Groups	Total Respond.#	Yes			No		
		N	%	C.I.(95%)	N	%	C.I.(95%)
TOTAL	3001	1986	62.7	(60.2-65.2)	1015	37.3	(34.8-39.8)
ADDs							
Barren River	211	132	60.9	(52.3-68.7)	79	39.1	(31.3-47.7)
Big Sandy	193	134	65.7	(57.7-72.9)	59	34.3	(27.1-42.3)
Bluegrass	228	144	62.0	(54.6-68.9)	84	38.0	(31.1-45.4)
Buffalo Trace	157	97	59.6	(50.7-67.9)	60	40.4	(32.1-49.3)
Cumberland Valley	201	150	68.2	(59.0-76.2)	51	31.8	(23.8-41.0)
FIVCO	229	161	70.2	(63.0-76.6)	68	29.8	(23.4-37.0)
Gateway	219	151	66.7	(58.8-73.8)	68	33.3	(26.2-41.2)
Green River	172	105	59.4	(50.9-67.3)	67	40.6	(32.7-49.1)
Kentucky River	217	163	75.5	(68.5-81.4)	54	24.5	(18.6-31.5)
KIPDA	189	116	59.7	(51.4-67.5)	73	40.3	(32.5-48.6)
Lake Cumberland	195	136	68.0	(60.2-75.0)	59	32.0	(25.0-39.8)
Lincoln Trail	193	122	58.7	(50.7-66.4)	71	41.3	(33.6-49.3)
Northern Kentucky	185	113	61.2	(53.3-68.6)	72	38.8	(31.4-46.7)
Pennyrile	208	133	62.3	(54.6-69.5)	75	37.7	(30.5-45.4)
Purchase	204	129	60.5	(52.7-67.8)	75	39.5	(32.2-47.3)

*This question is asked to respondents who reported having pain, aching, stiffness or swelling in or around a joint.

#Use caution in interpreting data for total respondents less than 50. N = Cell Size, NH = Non-Hispanic, % = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level).

Percentages are weighted to population characteristics.

Unless stated differently above, Missing, 'Don't know', and Refused responses are excluded.